



# Quasisymmetric Schur functions

Steph van Willigenburg  
University of British Columbia  
1984 Mathematics Rd  
Vancouver, BC.

AMS meeting May 2008

## Compositions and partitions

A **composition**  $\alpha_1 \dots \alpha_k$  of  $n$  is a list of positive integers whose sum is  $n$ : **2213**  $\vdash$  8.

A composition is a **partition** if  $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_k > 0$ : **3221**  $\vdash$  8.

Any composition **determines** a partition:  **$\lambda(2213) = 3221$** .

$\alpha = \alpha_1 \dots \alpha_k$  is a **coarsening** of  $\beta = \beta_1 \dots \beta_l$  ( $\beta$  is a **refinement** of  $\alpha$ ) if

$$\underbrace{\beta_1 + \dots + \beta_i}_{\alpha_1} \underbrace{\beta_{i+1} + \dots + \beta_j}_{\alpha_2} \dots \underbrace{\beta_m + \dots + \beta_l}_{\alpha_k}$$

is true: **53**  $\geq$  **2213**.

## Macdonald polynomials ...

1988: (Macdonald) Introduced  $\{P_\lambda(X; q, t)\}_\lambda$

$$(T) \quad P_\lambda(X; q, t) = s_\lambda(X) + \sum_{\mu < \lambda} s_\mu(X) \varepsilon_{\lambda, \mu}(q, t)$$

$$(\varepsilon_{\lambda, \mu}(q, t) \in \mathbb{Q}(q, t))$$

$$(O) \quad \langle P_\lambda(X; q, t), P_\mu(X; q, t) \rangle_{q, t} = 0 \quad \text{if } \lambda \neq \mu$$

... go nonsymmetric

$$P_\lambda(X; q, t) = \prod_{u \in \text{dg}'(\lambda^\circ)} (1 - q^{l(u)+1} t^{a(u)}) \sum_{\lambda(\mu)=\lambda} \frac{E_\gamma(X; q^{-1}, t^{-1})}{\prod (1 - q^{l(u)+1} t^{a(u)})}$$


letting  $q, t \rightarrow 0$

$$s_\lambda(X) = \sum_{\lambda(\mu)=\lambda} E_\gamma(X; \infty, \infty).$$

(Lascoux-Schützenberger (90); Reiner-Shimozono (95); Haglund, Haiman, Loehr (05); Mason (06))

## Composition diagrams and tableaux

The **composition diagram**  $\alpha = \alpha_1 \dots \alpha_k > 0$  is the array of **boxes** with  $\alpha_i$  boxes in row  $i$  from the **top**.



413

A **(standard) composition tableau** of **shape**  $\alpha$  is a filling of  $\alpha$  with (each first  $n$ )  $1, 2, 3, \dots$  such that

## Rules for composition tableaux I

- First column entries **strictly increase** top to bottom.
- Rows **weakly decrease** left to right.

*Example*

5	4	3	1
6			
8	7	2	

## Rules for composition tableaux II

- When the filling is **extended** for  $1 \leq i < j, 2 \leq k$

$$(j, k) \neq 0, (j, k) \geq (i, k) \Rightarrow (j, k) > (i, k - 1).$$

*Example*

5	4	3	1
6			
8	7	2	

## Rules for composition tableaux II

- When the filling is **extended** for  $1 \leq i < j, 2 \leq k$

$$(j, k) \neq 0, (j, k) \geq (i, k) \Rightarrow (j, k) > (i, k - 1).$$

*Example*

5	4	3	1
6	0	0	0
8	7	2	0



## Rules for composition tableaux II

- When the filling is **extended** for  $1 \leq i < j, 2 \leq k$

$$(j, k) \neq 0, (j, k) \geq (i, k) \Rightarrow (j, k) > (i, k - 1).$$

*Example*

5	4	3	1
6	0	0	0
8	7	2	0

## Rules for composition tableaux II

- When the filling is **extended** for  $1 \leq i < j, 2 \leq k$

$$(j, k) \neq 0, (j, k) \geq (i, k) \Rightarrow (j, k) > (i, k - 1).$$

*Example*

5	4	3	1
6	0	0	0
8	7	2	0

## Weights and weakness

Given a composition tableau  $T$  we have

$$x^T := x_1^{\#1s} x_2^{\#2s} x_3^{\#3s} \dots$$

*Example*

$$T = \begin{array}{|c|c|c|c|} \hline 5 & 4 & 3 & 1 \\ \hline 6 & & & \\ \hline 8 & 7 & 2 & \\ \hline \end{array} \quad x^T = x_1 x_2 x_3 x_4 x_5 x_6 x_7 x_8$$

A **weak composition**  $\gamma_1 \dots \gamma_k$  of  $n$  is a list of **non-negative** integers whose sum is  $n$ : **2021003**  $\vDash$  8.

## Demazure atoms

The **collapse** of a weak composition  $\gamma$ ,  $\alpha(\gamma)$ , is  $\gamma$  with 0's removed:  $\alpha(2021003) = 2213$ .

The **foundation** of a weak composition  $\gamma$ ,  $\mathcal{F}_o(\gamma)$ , is all  $i$  so  $\gamma_i > 0$ :  $\mathcal{F}_o(2021003) = \{1, 3, 4, 7\}$ .

Then

$$NS_\gamma(X) = E_{\gamma_n, \dots, \gamma_1}(x_n, \dots, x_1; \infty, \infty) = \sum_{\gamma} x^T$$

over composition tableaux shape  $\alpha(\gamma)$ , entries  $1, 2, 3, \dots, n$ , first column  $\mathcal{F}_o(\gamma)$ .

$NS_\gamma$  for  $\gamma = 1030$

1		
3	3	3

1		
3	3	2

1		
3	3	1

1		
3	2	2

1		
3	2	1

1		
3	1	1

$$NS_\gamma = x_1x_3^3 + x_1x_2x_3^2 + x_1^2x_3^2 + x_1x_2^2x_3 + x_1^2x_2x_3 + x_1^3x_3.$$

$NS_\gamma$  for  $\gamma = 1030$

1	0	0
3	3	3

1		
3	3	2

1		
3	3	1

1		
3	2	2

1		
3	2	1

1		
3	1	1

$$NS_\gamma = x_1x_3^3 + x_1x_2x_3^2 + x_1^2x_3^2 + x_1x_2^2x_3 + x_1^2x_2x_3 + x_1^3x_3.$$

$NS_\gamma$  for  $\gamma = 1030$

1		
3	3	3

1		
3	3	2

1		
3	3	1

1		
3	2	2

1		
3	2	1

1	0	0
3	1	1

$$NS_\gamma = x_1x_3^3 + x_1x_2x_3^2 + x_1^2x_3^2 + x_1x_2^2x_3 + x_1^2x_2x_3 + x_1^3x_3.$$

$NS_\gamma$  for  $\gamma = 1030$

1		
3	3	3

1		
3	3	2

1		
3	3	1

1		
3	2	2

1		
3	2	1

$$NS_\gamma = x_1x_3^3 + x_1x_2x_3^2 + x_1^2x_3^2 + x_1x_2^2x_3 + x_1^2x_2x_3 + \dots$$



## Quasisymmetric functions I

Let  $\mathcal{Q} \subset \mathbb{Q}[[x_1, x_2, \dots]]$  be the algebra of all **quasisymmetric functions**

$$\mathcal{Q} := \mathcal{Q}_0 \oplus \mathcal{Q}_1 \oplus \dots$$

where

$$\mathcal{Q}_n := \text{span}_{\mathbb{Q}}\{M_\alpha \mid \alpha = \alpha_1 \dots \alpha_k \models n\}$$

$$M_\alpha := \sum_{i_1 < i_2 < \dots < i_k} x_{i_1}^{\alpha_1} x_{i_2}^{\alpha_2} \dots x_{i_k}^{\alpha_k}.$$

*Example*  $M_{121} = \sum_{i_1 < i_2 < i_3} x_{i_1}^1 x_{i_2}^2 x_{i_3}^1 = x_1 x_2^2 x_3 + \dots$

## Quasisymmetric functions II

For  $\alpha \vDash n$  define

$$F_\alpha = \sum_{\beta \leq \alpha} M_\beta.$$

The  $F_\alpha$  are the **fundamental** quasisymmetric functions.

*Example*

$$F_{121} = M_{121} + M_{1111}$$

## Symmetric functions I

Let  $\Lambda \subset \mathbb{Q}[[x_1, x_2, \dots]]$  be the algebra of all **symmetric functions**

$$\Lambda := \Lambda_0 \oplus \Lambda_1 \oplus \dots$$

where

$$\Lambda_n := \text{span}_{\mathbb{Q}}\{m_\lambda \mid \lambda \vdash n\}$$


$$m_\lambda := \sum_{\lambda(\alpha)=\lambda} M_\alpha.$$

*Example*

$$m_{211} = M_{211} + M_{121} + M_{112}$$

## Contretableaux

The **diagram**  $\lambda = \lambda_1 \geq \dots \geq \lambda_k > 0$  is the array of **boxes** with  $\lambda_i$  boxes in row  $i$  from the **bottom**.



431

A **(standard) contretableau**  $T$  of **shape**  $\lambda$  is a filling of  $\lambda$  with (each first  $n$ )  $1, 2, 3, \dots$  so rows **weakly decrease** and columns **strictly increase**.

## Contretableaux

The **diagram**  $\lambda = \lambda_1 \geq \dots \geq \lambda_k > 0$  is the array of **boxes** with  $\lambda_i$  boxes in row  $i$  from the **bottom**.

5			
6	4	2	
8	7	3	1

431

A **(standard) contretableau**  $T$  of **shape**  $\lambda$  is a filling of  $\lambda$  with (each first  $n$ )  $1, 2, 3, \dots$  so rows **weakly decrease** and columns **strictly increase**.

## Descents and subsets

$D(T)$  is  $i$  where  $i + 1$  is strictly south and weakly east:

5			
6	4	2	
8	7	3	1

composition  $\alpha_1 \dots \alpha_k \vDash n \leftrightarrow$  subset  $\{i_1, \dots, i_{k-1}\} \subseteq [n - 1]$

$\beta$        $2312 \vDash 8 \leftrightarrow \{2, 5, 6\} \subseteq [7]$        $S(\beta)$

## Symmetric functions II

For  $\lambda \vdash n$  define

$$s_\lambda = \sum_{\beta} d_{\lambda\beta} F_\beta$$

where  $d_{\lambda\beta}$  = number of standard tableaux  $T$  of shape  $\lambda$  and  $D(T) = S(\beta)$ .

The  $s_\lambda$  are the **Schur** functions.

*Example*

$$s_{21} = F_{21} + F_{12}$$

from

2	
3	1

1	
3	2

## Quasisymmetric Schur functions

If  $m_\lambda = \sum_{\lambda(\alpha)=\lambda} M_\alpha$  then  $s_\lambda = \sum_{\lambda(\alpha)=\lambda} \mathcal{S}_\alpha$

but what is  $\mathcal{S}_\alpha$ ?

The quasisymmetric Schur function is given by ...



## Quasisymmetric Schur functions

If  $m_\lambda = \sum_{\lambda(\alpha)=\lambda} M_\alpha$  then  $s_\lambda = \sum_{\lambda(\alpha)=\lambda} \mathcal{S}_\alpha$

but what is  $\mathcal{S}_\alpha$ ?

The quasisymmetric Schur function is given by

$$\mathcal{S}_\alpha = \sum_{\alpha(\gamma)=\alpha} N S_\gamma.$$

$S_\alpha$  for  $\alpha = 12$

1	
2	2

1	
3	2

1	
3	3

2	
3	3

$$\begin{aligned} S_{12} &= NS_{120} + NS_{102} + NS_{012} \\ &= x_1x_2^2 + x_1x_2x_3 + x_1x_3^2 + x_2x_3^2 \\ &= M_{12} + M_{111} \end{aligned}$$

and

$$s_{21} = F_{21} + F_{12} = M_{21} + M_{12} + 2M_{111} = s_{21} + S_{12}.$$

## What properties should $\mathcal{S}_\alpha$ have?

- $\mathbb{Z}$ -basis for  $\mathcal{Q}$ .
- Expression in  $F_\beta$ .
- Quasisymmetric Kostka numbers.
- Quasisymmetric Pieri rules.
- Quasisymmetric LR rule.

## What properties should $\mathcal{S}_\alpha$ have?

- $\mathbb{Z}$ -basis for  $\mathcal{Q}$ .
- Expression in  $F_\beta$ .
- Quasisymmetric Kostka numbers.
- Quasisymmetric Pieri rules.
- Quasisymmetric LR rule.

## Expression in $F_\beta$

$\mathcal{D}(T)$  is  $i$  where  $i + 1$  is north in column (immediately south if both in first column), or strictly east:

4	3	2
5	1	

For  $\lambda \vdash n$

$$s_\lambda = \sum_{\beta} d_{\lambda\beta} F_\beta$$

where  $d_{\lambda\beta}$  = number of standard tableaux  $T$  of shape  $\lambda$  and  $D(T) = S(\beta)$ .

## Expression in $F_\beta$

$\mathcal{D}(T)$  is  $i$  where  $i + 1$  is north in column (immediately south if both in first column), or strictly east:

4	3	2
5	1	

For  $\alpha \vDash n$

$$S_\alpha = \sum_{\beta} d_{\alpha\beta} F_\beta$$

where  $d_{\alpha\beta}$  = number of standard tableaux  $T$  of shape  $\alpha$  and  $\mathcal{D}(T) = S(\beta)$ .

$S_\alpha$  for  $\alpha = 32$

3	2	1
5	4	

4	3	1
5	2	

4	3	2
5	1	

and


$$S_{32} = F_{32} + F_{221} + F_{131}.$$

## Skew diagrams and strips


For  $\lambda, \mu$  the skew diagram  $\lambda/\mu$  of size  $|\lambda/\mu|$  is the array of boxes contained in  $\lambda$  but not in  $\mu$ .

A skew diagram  $\lambda/\mu$  is a row (column) strip if

no 2 boxes in same column (row).



row strip



col strip




## Skew diagrams and strips


For  $\lambda, \mu$  the skew diagram  $\lambda/\mu$  of size  $|\lambda/\mu|$  is the array of boxes contained in  $\lambda$  but not in  $\mu$ .

A skew diagram  $\lambda/\mu$  is a row (column) strip if

no 2 boxes in same column (row).



row strip



col strip

## Pieri rules for partitions $\lambda, \mu$

$$s_n s_\lambda = \sum_{\mu} s_\mu \quad s_{1^n} s_\lambda = \sum_{\mu} s_\mu$$

- $\delta = \mu/\lambda$  row strip,  $|\delta| = n$ .
- $\varepsilon = \mu/\lambda$  column strip,  $|\varepsilon| = n$ .

## Removal from compositions

Let  $\alpha = \alpha_1 \dots \alpha_k$  have largest part  $m$ , and  $s \in [m]$

$$rem_s(\alpha) = \alpha_1 \dots \alpha_{i-1} (s-1) \alpha_{i+1} \dots \alpha_k$$

for largest  $i$ .

If  $S = \{s_1 < \dots < s_j\}$  then

$$row_S(\alpha) = rem_{s_1}(\dots (rem_{s_{j-1}}(rem_{s_j}(\alpha))) \dots).$$

If  $M = \{m_1 \leq \dots \leq m_j\}$  then

$$col_M(\alpha) = rem_{m_j}(\dots (rem_{m_2}(rem_{m_1}(\alpha))) \dots).$$

$rem_s$  in action

$rem_s$  removes rightmost box from lowest row length  $s$ .

*Example*

$$rem_1(113) = \begin{array}{|c|} \hline \square \\ \hline \bullet \\ \hline \square & \square & \square \\ \hline \end{array} = \begin{array}{|c|} \hline \square \\ \hline \square & \square & \square \\ \hline \end{array} = 13$$

## Quasisymmetric Pieri rules for compositions $\alpha, \beta$

$$\mathcal{S}_n \mathcal{S}_\alpha = \sum_{\beta} \mathcal{S}_\beta \quad \mathcal{S}_{1^n} \mathcal{S}_\alpha = \sum_{\beta} \mathcal{S}_\beta$$

- $\delta = \lambda(\beta)/\lambda(\alpha)$  row strip,  $|\delta| = n$ .
- $row_{\mathcal{S}(\delta)}(\beta) = \alpha$ .
- $\varepsilon = \lambda(\beta)/\lambda(\alpha)$  column strip,  $|\varepsilon| = n$ .
- $col_{M(\varepsilon)}(\beta) = \alpha$ .

## $\mathcal{S}_1\mathcal{S}_{13}$ in action

41/31, 32/31, 311/31, 311/31

with row strips containing a box in column 4, 2, 1, 1 respectively.

$$\begin{aligned} \text{row}_{\{4\}}(14) &= \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \bullet \\ \hline \end{array} & \text{row}_{\{2\}}(23) &= \begin{array}{|c|c|c|} \hline & \bullet & \\ \hline & & \\ \hline \end{array} \\ \text{row}_{\{1\}}(131) &= \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline \bullet & & \\ \hline \end{array} & \text{row}_{\{1\}}(113) &= \begin{array}{|c|c|c|} \hline & & \\ \hline \bullet & & \\ \hline & & \\ \hline \end{array} \end{aligned}$$

and

$$\mathcal{S}_1\mathcal{S}_{13} = \mathcal{S}_{14} + \mathcal{S}_{23} + \mathcal{S}_{131} + \mathcal{S}_{113}.$$