


HOMEWORK #3 - MATH 4160

ASSIGNED: OCT 30, 2014 DUE: NOV 11, 2013

Write your homework solutions neatly and clearly (type!). Provide full explanations and justify all of your answers.

- (1) Give an argument which explains why the number of ways of coloring the vertices of the following graph with k colors such that no two vertices connected with an edge is colored the same, is equal to $k^5 - 7k^4 + 19k^3 - 23k^2 + 10k$ (the number of ways of coloring this graph with no restriction on the colors is k^5). Phrase your explanation in terms of an applications of the addition principle and the multiplication principle.



- (2) Find the generating function for the number of words of length n using letters $a, b, c, d, e, f, g, h, i$ such that
- all 9 letters occur without restriction
 - at least one of the first 6 letters appears
 - the first 6 letters each appear at least once and the last three appear an even number of times
 - the first 6 letters each appear at least once and the last six appear an even number of times
 - at least one of the first 6 letters appears and the total number of the last 6 letters is even
- Use your generating function to find the number of number of words of length 20 with the restrictions above.
- (3) In class we found the exponential generating function for the Bell numbers $B(n)$ which are defined by the recurrence $B(0) = 1$, $B(1) = 1$ and $B(n + 1) = \sum_{i=1}^n \binom{n}{i} B(n - i)$ for all $n \geq 1$. We found that $B(x) = \sum_{n \geq 0} B(n) \frac{x^n}{n!} = e^{e^x - 1}$. Recall that the Stirling numbers of the second kind $S_{n,k}$ are defined as the number of set partitions into k parts. They are defined recursively as $S_{0,0} = 1$, $S_{n,1} = S_{n,n} = 1$ for all $n \geq 1$, and $S_{n,k} = 0$ if $k > n$. Moreover $S_{n+1,k} = kS_{n,k} + S_{n,k-1}$ for $n \geq 0$ and $1 \leq k \leq n$.

Refine the computation that gives the formula for $B(x) = \sum_{n \geq 0} B(n) \frac{x^n}{n!} = e^{e^x - 1}$ to show that

$$S(x, q) = \sum_{n \geq 0} \sum_{k \geq 0} S_{n,k} q^k \frac{x^n}{n!} = e^{q(e^x - 1)}.$$

- (4) Use the result of the previous problem to give the generating function for the number of set partitions into an odd number of parts. That is, if we let

$$B_o(n) = \sum_{k=0}^{\lceil n/2 \rceil - 1} S_{n, 2k+1},$$

then find a formula for $B_o(x) = \sum_{n \geq 0} B_o(n) \frac{x^n}{n!}$.

- (5) Given the generating function $A(x) = \sum_{n \geq 0} a_n x^n = a_0 + a_1 x + a_2 x^2 + \dots$, find a formula for the generating function

$$\tilde{A}(x) = a_1 + a_0 x + a_3 x^2 + a_2 x^3 + a_5 x^4 + a_4 x^5 + \dots$$

If $B(x) = \sum_{n \geq 0} b_n \frac{x^n}{n!} = b_0 + b_1 \frac{x}{1!} + b_2 \frac{x^2}{2!} + \dots$ is an exponential generating function, find a formula for the generating function

$$\tilde{B}(x) = b_1 + b_0 \frac{x}{1!} + b_3 \frac{x^2}{2!} + b_2 \frac{x^3}{3!} + b_5 \frac{x^4}{4!} + b_4 \frac{x^5}{5!} + \dots$$