

Combinatorial Hopf Algebras.

YORK



UNIVERSITÉ

UNIVERSITY

Nantel Bergeron

York Research Chair in Applied Algebra

www.math.yorku.ca/bergeron

[with J.Y. Thibon ... and many more]





Outline


- What would be a good **gift** for a mathematician?
- What is a **Combinatorial Hopf Algebra**?
- **Sym** is a **strong, realizable** CHA with **character**.
- On strong CHA (**categorification**)
- On realizable CHA (**word combinatorics** and **quotients**).

Combinatorial Hopf Algebra

$H = \bigoplus_{n \geq 0} H_n$ a graded connected Hopf algebra is **CHA** if

- (weak) There is a distinguished (combinatorial) basis with positive integral structure coefficients (from **Hopf monoid**).
- (strong) The structure is obtained from representation operation (from **categorification**).
- (real.) It can be realized in a space of series in variables. (it is **realizable**)
- (char.) It has a distinguished character. (with **character**)

Combinatorial Hopf Algebra



Sym is the model CHA

Sym is the space of symmetric functions $\mathbb{Z}[h_1, h_2, \dots]$, with $\deg(h_k) = k$ and

$$\Delta(h_k) = \sum_{i=0}^k h_i \otimes h_{k-i}.$$

Sym is the model CHA

Sym

Sym is the model CHA

Sym is the space of symmetric functions $\mathbb{Z}[h_1, h_2, \dots]$, with $\deg(h_k) = k$ and

$$\Delta(h_k) = \sum_{i=0}^k h_i \otimes h_{k-i}.$$

It is the functorial image of a **Hopf Monoid Π** :

For any finite set J let $\Pi[J] = \{A : A \vdash J\}$ the **set partitions** of J .


Product and **Coproduct**:

combinatorial constructions on set partitions

It correspond to **flats** of the hyperplane arrangement of type A .

Sym is the model CHA

Hopf Monoid Π



Hopf structure on $\bigoplus_{n \geq 0} K_0(S_n)$

$K_0(S) = \bigoplus_{n \geq 0} K_0(S_n)$ is the space of S_n -modules up to isomorphism

- **Basis:** Irreducible modules S^λ
- **Structure:**

$$M * N = \text{Ind}_{S_n \times S_m}^{S_{n+m}} M \otimes N$$

$$\Delta M = \bigoplus_{k=0}^n \text{Res}_{S_k \times S_{n-k}}^{S_n} M$$

- $\mathcal{F}: K_0(S) \rightarrow \text{Sym}$ is an **isomorphism** of graded Hopf algebra where $\mathcal{F}(S^\lambda) = s_\lambda$

Sym is the model CHA

Hopf Monoid Π

$\{A\}_{A \vdash J}$

\overline{K}

Categorification

$\{S^\lambda\}_{\lambda \vdash n}$

$\{h_\lambda\}_{\lambda \vdash n}$
 $\{m_\lambda\}_{\lambda \vdash n}$

Sym

$\{s_\lambda\}_{\lambda \vdash n}$

\mathcal{F}

Realization of Sym

$$Sym \hookrightarrow \lim_{n \rightarrow \infty} \mathbb{Q}[x_1, x_2, \dots, x_n]$$

Allows us to understand coproducts, internal coproduct, plethysm, Cauchy kernel, ...

Sym is the model CHA

Hopf Monoid Π

$\{A\}_{A \vdash J}$

\overline{K}

Categorification

$\{S^\lambda\}_{\lambda \vdash n}$

\mathcal{F}

$\{h_\lambda\}_{\lambda \vdash n}$
 $\{m_\lambda\}_{\lambda \vdash n}$

Sym

$\{s_\lambda\}_{\lambda \vdash n}$

$\lim_{n \rightarrow \infty} \mathbb{Q}[x_1, x_2, \dots, x_n]$

Sym with a Hopf Character

$$\begin{aligned}\zeta_0: \quad Sym &\longrightarrow \mathbb{Q} \\ f(x_1, x_2, \dots) &\mapsto f(1, 0, \dots)\end{aligned}$$

(Sym, ζ_0) is a terminal object for (H, ζ) cocommutative:

$$\begin{array}{ccc} H & \cdots \longrightarrow & Sym \\ \zeta \searrow & & \swarrow \zeta_0 \\ & \mathbb{Q} & \end{array}$$


$$\zeta_0^* = \sum_{n \geq 0} h_n$$

$$\Omega(X) = \sum_{n \geq 0} h_n(X) = \prod_{x \in X} \frac{1}{1 - x}$$

Sym is the model CHA

Hopf Monoid Π

Categorification



Toward Categorification

Consider a graded algebra $A = \bigoplus_{n \geq 0} A_n$

- Each A_n is an algebra.
- $\dim A_0 = 1$ and $\dim A_n < \infty$.
- $\rho_{n,m}: A_n \otimes A_m \hookrightarrow A_{n+m}$; injective algebra homomorphism
- A_{n+m} is projective bilateral submodule of $A_m \otimes A_m$.
- Right and left projective structure of A_{n+m} are compatible.
- There is a Mackey formula linking induction and restriction

A is a tower of algebra

Toward Categorification

Consider a tower of algebras $A = \bigoplus_{n \geq 0} A_n$

Let $K_0(A) = \bigoplus_{n \geq 0} K_0(A_n)$ is the space of (projective) A_n -modules up to isomorphism and modulo short exact sequences

- $K_0(A)$ is a **graded Hopf algebra**:

$$M * N = \text{Ind}_{A_n \otimes A_m}^{A_{n+m}} M \otimes N$$

$$\Delta M = \bigoplus_{k=0}^n \text{Res}_{A_k \otimes A_{n-k}}^{A_n} M$$

- H is a **strong CHA** if there is an **isomorphism**

$$\mathcal{F}: K_0(A) \rightarrow H$$

Example of Tower of Algebras

$$\mathbb{Q}S = \bigoplus_{n \geq 0} \mathbb{Q}S_n:$$

$$\mathcal{F}: K_0(\mathbb{Q}S) \rightarrow \textcolor{red}{Sym}$$

$$H(0) = \bigoplus_{n \geq 0} H_n(0): \textcolor{green}{[Krob-Thibon]}$$

$$\mathcal{F}: K_0(H(0)) \rightarrow \textcolor{red}{NSym}$$

$$\mathcal{F}: G_0(H(0)) \rightarrow \textcolor{red}{QSym}$$

$$HC(0) = \bigoplus_{n \geq 0} HC_n(0): \textcolor{green}{[B-Hivert-Thibon]} \dots \text{Peak algebras} \dots$$

seams rare?

Obstruction to Tower of algebras?

Consider a tower of algebras $A = \bigoplus_{n \geq 0} A_n$

where $K_0(A)$ and $G_0(A)$ are **graded dual Hopf algebra**:

THEOREM[B-Lam-Li]

if A is a tower of algebras, then $\dim(A_n) = r^n n!$

this is very restrictive...

Tower of Supercharacters [... B ... Novelli ... Thibon ...]

- Unipotent upper triangular matrices over finite Fields \mathbf{F}_q : $U_n(q)$.
- **Superclasses** in $U_n(q)$: $A \cong B \iff (A - I) = M(B - I)N$
- **Supercharacters** χ : characters constant on superclasses:

$$\Delta(\chi) = \sum_{A+B=[n]} \text{Res}_{U_{|A|}(q) \times U_{|B|}(q)}^{U_n(q)} \chi$$

$$\chi \cdot \psi = \text{Inf}_{U_n(q) \times U_m(q)}^{U_{n+m}(q)} \chi \otimes \psi = (\chi \otimes \psi) \circ \pi$$

where $\pi: U_{n+m}(q) \rightarrow U_n(q) \times U_m(q)$.

- $\mathcal{F}: K_0\left(\bigoplus_{n \geq 0} U_n(2)\right) \rightarrow \text{NC Sym}$ is iso.

NC Sym symmetric functions in non-commutative variables.

Some open questions

(Q-1) Find **other examples** of Categorification (Can we do *NCQsym* (quasi-symmetric in non commutative variables)?

(Q-2) **Tower of algebra** A (axiomatization with superclasses/
supermodules and Harish-Chandra induction:

$$\text{Ind} \circ \text{Inf} \quad \text{and} \quad \text{Def} \circ \text{Res} \quad).$$

About Realization

Many CHA are realized: Sym, NSym, QSym, NCSym, •••

Can we described all

$$H \hookrightarrow \mathbb{Q}\langle x_1, x_2, \dots \rangle$$

with monomial basis (equivalence classes on words) [Giraldo].

[B-Hohlweg] Monomial basis embeddings

$$H \hookrightarrow SSym$$


(Q-3) Realization Theory: Can we describe monomial embeddings

$$H \hookrightarrow \mathbb{Q}M$$

for different monoid M

Realization Quotients

Hopf algebras



Reverse Lex and Gröbner basis

$$\mathbb{Q}[x_1, \dots, x_{n+1}] \xrightarrow{x_n=0} \mathbb{Q}[x_1, \dots, x_n]$$

$$H[x_1, \dots, x_{n+1}] \xrightarrow{x_n=0} H[x_1, \dots, x_n]$$

G_n G-basis of ideal $\langle H[x_1, \dots, x_n]^+ \rangle$:

$$G_{n+1} \xrightarrow{x_n=0} G_n$$

$$g(x_1, \dots, x_{n+1}) \mapsto \begin{cases} 0 & \text{if } LT(g)|_{x_n=0} = 0 \\ \tilde{g} & \text{if } LT(g)|_{x_n=0} = LT(\tilde{g}) \neq 0 \end{cases}$$

B_n basis of quotient $\mathbb{Q}[x_1, \dots, x_n] / \langle H[x_1, \dots, x_n]^+ \rangle$:

$$B_{n+1} \longleftarrow B_n$$

Reverse Lex and Gröbner basis

$$\mathbb{Q}[x_1, \dots, x_{n+1}] \xrightarrow{x_n=0} \mathbb{Q}[x_1, \dots, x_n]$$

$$H[x_1, \dots, x_{n+1}] \xrightarrow{x_n=0} H[x_1, \dots, x_n]$$

$$G_{n+1} \xrightarrow{x_n=0} G_n$$

$$g(x_1, \dots, x_{n+1}) \mapsto \begin{cases} 0 & \text{if } LT(g)|_{x_n=0} = 0 \\ \tilde{g} & \text{if } LT(g)|_{x_n=0} = LT(\tilde{g}) \neq 0 \end{cases}$$

$$B_{n+1} \xleftarrow{\quad} B_n$$


mult by x_n

mult by x_n^2

mult by x_n^3

...

Realization Quotients



2
Jan 2005


Realization Quotients

Diagonally S_n -covariants

Haiman and others...

$(n+1)^{n-1}$


Diagonally Temperley Lieb
covariants Aval Bergeron Bergeron



3

Jan 2005

Realization Quotients



Diagonally TL-covariants

[Aval Bergeron Bergeron]

$$D_n := \mathbb{Q}[x_1, x_2, \dots, x_n; y_1, \dots, y_n] / \langle \text{DQSym}^+ \rangle$$

Conjectured bigraded Hilbert series:

$$\dim_{qt} D_1 = \begin{bmatrix} 1 \end{bmatrix} \quad \dim_{qt} D_2 = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$

$$\dim_{qt} D_3 = \begin{bmatrix} 2 & 0 & 0 \\ 2 & 2 & 0 \\ 1 & 2 & 2 \end{bmatrix} \quad \dim_{qt} D_4 = \begin{bmatrix} 5 & 0 & 0 & 0 \\ 5 & 5 & 0 & 0 \\ 3 & 7 & 5 & 0 \\ 1 & 3 & 5 & 5 \end{bmatrix}$$

degree in q \uparrow $n-1$

$$\dim_{qt} D_5 = \begin{bmatrix} 14 & 0 & 0 & 0 & 0 \\ 14 & 14 & 0 & 0 & 0 \\ 9 & 24 & 14 & 0 & 0 \\ 4 & 14 & 24 & 14 & 0 \\ 1 & 4 & 9 & 14 & 14 \end{bmatrix}$$

0 \rightarrow degree in t $n-1$


5
Jan 2005

Diagonally TL-covariants

[Aval Bergeron Bergeron]

$$D_n := \mathbb{Q}[x_1, x_2, \dots, x_n; y_1, \dots, y_n] / \langle \text{DQSym}^+ \rangle$$

Conjectured explicit monomial basis:
for example to build for $n=4$ and bidegree $(1,1)$



6
Jan 2005



About family of Realization

(Q-4) Prove previous question about Hilbert series

(Q-5) Realized Quotient in general

...

M E R C I



T H A N K S

G R A C I A S

